

Action-angle coordinates around a noncompact invariant manifold of a completely integrable Hamiltonian system

E.FIORANI¹

Dipartimento di Matematica "F.Enriques", Università di Milano, 20133, Milano, Italy

G.GIACHETTA²,

Dipartimento di Matematica e Fisica, Università di Camerino, 62032 Camerino (MC), Italy

G. SARDANASHVILY³

Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

Abstract. A trivial bundle of regular connected invariant manifolds of a completely integrable Hamiltonian system can be provided with action-angle coordinates.

MSC (2000): 37J35, 70H06

By virtue of the classical theorem [1, 2], an autonomous completely integrable Hamiltonian system admits action-angle coordinates around a connected regular compact invariant manifold. We show that there is a system of action-angle coordinates on an open neighbourhood U of a connected regular invariant manifold M if Hamiltonian vector fields of first integrals on U are complete and the foliation of U by invariant manifolds is trivial. If M is a compact regular invariant manifold, these conditions always hold.

Theorem 1. Let M be a connected invariant manifold of a completely integrable Hamiltonian system $\{F_\lambda\}$, $\lambda = 1, \dots, n$, on a symplectic manifold (Z, Ω) . Let U be an open neighbourhood of M such that: (i) $\{F_\lambda\}$ have no critical points in U , (ii) the Hamiltonian vector fields of the first integrals F_λ on U are complete, and (iii) the submersion $\times F_\lambda : U \rightarrow \mathbb{R}^n$ is a trivial bundle of invariant manifolds over a domain $V \subset \mathbb{R}^n$. Then U is isomorphic to the symplectic annulus

$$W = V \times (\mathbb{R}^{n-m} \times T^m), \quad (1)$$

provided with the action-angle coordinates

$$(I_1, \dots, I_n; x^1, \dots, x^{n-m}; \phi^1, \dots, \phi^m)$$

such that the symplectic form on W reads

$$\Omega = dI_a \wedge dx^a + dI_i \wedge d\phi^i,$$

¹E-mail: fiorani@mat.unimi.it

²E-mail: giovanni.giachetta@unicam.it

³E-mail: sard@grav.phys.msu.su; URL: <http://webcenter.ru/~sardan/>

and the first integrals F_λ depend only on the action coordinates (I_α) .

Proof. In accordance with the well-known theorem [1], the invariant manifold M is diffeomorphic to the product $\mathbb{R}^{n-m} \times T^m$, which is the group space of the quotient $G = \mathbb{R}^n / \mathbb{Z}^m$ of the group $\mathbb{R}^n$ generated by Hamiltonian vector fields ϑ_λ of first integrals F_λ on M . Namely, M is provided with the group space coordinates $(y^\lambda) = (s^a, \varphi^i)$ where φ^i are linear functions of parameters s^λ along integral curves of the Hamiltonian vector fields ϑ_λ on U . Let (J_λ) be coordinates on V which are values of first integrals F_λ . Let us choose a trivialization of the fiber bundle $U \rightarrow V$ seen as a principal bundle with the structure group G . We fix its global section χ . Since parameters s^λ are given up to a shift, let us provide each fiber M_J , $J \in V$, with the group space coordinates (y^λ) centered at the point $\chi(J)$. Then $(J_\lambda; y^\lambda)$ are bundle coordinates on the annulus W (1). Since M_J are Lagrangian manifolds, the symplectic form Ω on W is given relative to the bundle coordinates $(J_\lambda; y^\lambda)$ by the expression

$$\Omega = \Omega^{\alpha\beta} dJ_\alpha \wedge dJ_\beta + \Omega_\beta^\alpha dy^\beta. \quad (2)$$

By the very definition of coordinates (y^λ) , the Hamiltonian vector fields ϑ_λ of first integrals take the coordinate form $\vartheta_\lambda = \vartheta_\lambda^\alpha (J_\mu) \partial_\alpha$. Moreover, since the cyclic group S^1 can not act transitively on $\mathbb{R}$, we have

$$\vartheta_a = \partial_a + \vartheta_a^i (J_\lambda) \partial_i, \quad \vartheta_i = \vartheta_i^k (J_\lambda) \partial_k. \quad (3)$$

The Hamiltonian vector fields ϑ_λ obey the relations

$$\vartheta_\lambda \lrcorner \Omega = -dJ_\lambda, \quad \Omega_\beta^\alpha \vartheta_\lambda^\beta = \delta_\lambda^\alpha. \quad (4)$$

It follows that Ω_β^α is a nondegenerate matrix and $\vartheta_\lambda^\alpha = (\Omega^{-1})_\lambda^\alpha$, i.e., the functions Ω_β^α depend only on coordinates J_λ . A substitution of (3) into (4) results in the equalities

$$\Omega_b^a = \delta_b^a, \quad \vartheta_a^\lambda \Omega_\lambda^i = 0, \quad (5)$$

$$\vartheta_i^k \Omega_k^j = \delta_i^j, \quad \vartheta_i^k \Omega_k^a = 0. \quad (6)$$

The first of the equalities (6) shows that the matrix Ω_k^j is nondegenerate, and so is the matrix ϑ_i^k . Then the second one gives $\Omega_k^a = 0$.

By virtue of the well-known Künneth formula for the de Rham cohomology of a product of manifolds, the closed form Ω (2) on W (1) is exact, i.e., $\Omega = d\Xi$ where Ξ reads

$$\Xi = \Xi^\alpha (J_\lambda, y^\lambda) dJ_\alpha + \Xi_i (J_\lambda) d\varphi^i + \partial_\alpha \Phi (J_\lambda, y^\lambda) dy^\alpha,$$

where Φ is a function on W . Taken up to an exact form, Ξ is brought into the form

$$\Xi = \Xi'^\alpha (J_\lambda, y^\lambda) dJ_\alpha + \Xi_i (J_\lambda) d\varphi^i. \quad (7)$$

Owing to the fact that components of $d\Xi = \Omega$ are independent of y^λ and obey the equalities (5) – (6), we obtain the following.

(i) $\Omega_i^a = -\partial_i \Xi'^a + \partial^a \Xi_i = 0$. It follows that $\partial_i \Xi'^a$ is independent of φ^i , i.e., Ξ'^a is affine in φ^i and, consequently, is independent of φ^i since φ^i are cyclic coordinate. Hence, $\partial^a \Xi_i = 0$, i.e., Ξ_i is a function only of coordinates J_j .

(ii) $\Omega_i^k = -\partial_i \Xi'^k + \partial^k \Xi_i$. Similarly to item (i), one shows that Ξ'^k is independent of φ^i and $\Omega_i^k = \partial^k \Xi_i$, i.e., $\partial^k \Xi_i$ is a nondegenerate matrix.

(iii) $\Omega_b^a = -\partial_b \Xi'^a = \delta_b^a$. Hence, $\Xi'^a = -s^a + D^a(J_\lambda)$.

(iv) $\Omega_b^i = -\partial_b \Xi'^i$, i.e., Ξ'^i is affine in s^a .

In view of items (i) – (iv), the Liouville form Ξ (7) reads

$$\Xi = x^a dJ_a + [D^i(J_\lambda) + B_a^i(J_\lambda) s^a] dJ_i + \Xi_i(J_j) d\varphi^i,$$

where we put

$$x^a = -\Xi'^a = s^a - D^a(J_\lambda). \quad (8)$$

Since the matrix $\partial^k \Xi_i$ is nondegenerate, one can introduce new coordinates $I_i = \Xi_i(J_j)$, $I_a = J_a$. Then we have

$$\Xi = -x^a dI_a + [D'^i(I_\lambda) + B_a^i(I_\lambda) s^a] dI_i + I_i d\varphi^i.$$

Finally, put

$$\phi^i = \varphi^i - [D'^i(I_\lambda) + B_a^i(I_\lambda) s^a] \quad (9)$$

in order to obtain the desired action-angle coordinates

$$I_a = J_a, \quad I_i(J_j), \quad x^a(J_\lambda, s^a), \quad \phi^i(J_\lambda, y^\lambda).$$

These are bundle coordinates on $U \rightarrow V$ where the coordinate shifts (8) – (9) correspond to a choice of another trivialization of $U \rightarrow V$. QED

In particular, Theorem 1 enables one to introduce action-angle coordinates for time-dependent completely integrable Hamiltonian systems whose invariant manifolds are never compact because of the time axis [3]. Another application of Theorem 1 is Jacobi fields of a completely integrable system. They also make up a completely integrable system whose invariant manifolds are never compact [4].

References

- [1] Arnold V (Ed.) 1988 *Dynamical Systems III* (Berlin: Springer-Verlag)
- [2] Lazutkin V 1993 *KAM Theory and Semiclassical Approximations to Eigenfunctions* (Berlin: Springer-Verlag)
- [3] Giachetta G, Mangiarotti L and Sardanashvily G, E-print arXiv.org: math.DS/0204151
- [4] G.Giachetta, L.Mangiarotti and G.Sardanashvily, E-print arXiv.org: math-ph/0205026